

Southern Infrared Proper Motion Survey I: Discovery of New High Proper Motion Stars From First Full Hemisphere Scan

N.R. Deacon, N. C. Hambly and J. A. Cooke

Institute for Astronomy, University of Edinburgh, Blackford Hill, Edinburgh EH9 3HJ
e-mail: nd@roe.ac.uk, n.hambly@roe.ac.uk, j.cooke@roe.ac.uk

Received —; accepted —

Abstract. We present the first results from the Southern Infrared Proper Motion Survey. Using 2 Micron All Sky Survey data along with that of the SuperCOSMOS sky survey we have been able to produce the first widefield infrared proper motion survey. Having targeted the survey to identify nearby M, L and T dwarfs we have discovered 72 such new objects with proper motions greater than $0.5''/\text{yr}$ with 10 of these having proper motions in excess of $1''/\text{yr}$. The most interesting of these objects is SIPS1259-4336 a late M dwarf. We have calculated a trigonometric parallax for this object of $\pi = 276 \pm 41$ milliarcseconds yielding a distance of $3.62 \pm 0.54 \text{ pc}$. We have also discovered a common proper motion triple system and an object with a common proper motion with LHS 128. The survey completeness is limited by the small epoch differences between many 2MASS and UKI observations. Hence we only recover 22% of Luyten objects with favourable photometry. However the Luyten study is itself unquantifiably incomplete. We discuss the prospect of enhancing the survey volume by reducing the lower proper motion limit.

Key words. Astrometry – Stars: Low mass, brown dwarfs

1. Introduction

The Solar Neighbourhood provides an opportunity to constrain the field mass and luminosity functions of Low Mass Stars and Brown Dwarfs. As Low Mass Stars and Brown Dwarfs represent the products of star formation they provide clues as to the processes involved. They also represent a sink for baryonic matter and hence may explain a small proportion of Galactic dark matter. In addition nearby multiples can give clues about the formation of such systems.

Proper motion surveys are a useful tool for discovering nearby stars. The benchmark for such surveys is Luyten's Half Arcsecond Catalogue (hereafter LHS) (1979) containing 3587 stars with $\mu \geq 0.5''/\text{yr}$. This survey is incomplete in the southern sky and excludes fields in the galactic plane. Modern high proper motion searches seek to complete Luyten's work in the southern sky and to identify objects Luyten failed to find. Some of these use manual blinking techniques such as the Calan-ESO proper motion catalogue (Ruiz et al 2001) and the study of Wroblewski and Costa (2001). Recent computational methods include those of Pokorny et al (2003) and Scholz et al (2002). Pokorny lists 6206 stars with $\mu \geq 0.18''/\text{yr}$ found using SuperCOSMOS scans of 131 Schmidt fields. Scholz found 15 stars with high proper motions from APM scans of UKST plates. Eight of these have distances less than 25 pc. Lepine and Shara (2002) have used POSS

plates to search in the northern hemisphere. This has been done with the SUPERBLINK algorithm, in which plates from two different epochs (POSS I and POSS II) are aligned and the later, better POSS II plates degraded to make them of comparable quality to the POSS I plates. The two images are then subtracted making the moving, high proper motion objects obvious. Teegarden et al's (2003) recent discovery was made using data from the SkyMorph search for near earth objects, which also picks up stars with a very high proper motion. Most recently Hambly et al. (2004) have used SuperCOSMOS data to identify five new high proper motion stars south of $\delta = -60^\circ$.

However the majority of these surveys utilise optical R plate data and many nearby stars are so red that they are barely visible on such plates. Low Mass Stars and Brown Dwarfs are bright in the near infrared I , J , H and K_s . This means that a proper motion search in the near infrared is ideal for identifying such objects. Near infrared proper motion surveys have already been used to discover low mass companions to nearby stars (Scholz et al., 2004) and to confirm that companions are co-moving with the primary (Seifahrt et al., 2004). However these are targeted at known nearby stars and are not useful for discovering new systems. Many proper motion surveys are also incomplete. SCR1845–6357 for example was missed by all surveys prior to Hambly et al (2004) despite being rela-

tively bright at $R \approx 16$ and having a high proper motion $\mu > 2''/\text{yr}$. It is likely that similar objects are still to be discovered. In order to identify late M, L and T dwarfs a proper motion survey in the near infrared is required. For our survey we used SuperCOSMOS I data and data in J, H, K_s from the 2MASS Point Source Catalogue as our two epochs. 2MASS objects likely to be nearby low mass stars or brown dwarfs were selected. These were then paired with likely I plate counterparts. This constitutes the first widefield infrared proper motion survey ideal for identifying nearby Low Mass Stars and Brown Dwarfs. Here we present details of the method used as well as a list of objects found with $\mu > 0.5''/\text{yr}$.

2. Candidate Identification

2.1. Initial 2MASS object selection

Firstly candidate 2MASS images had to be selected. To ensure that these objects were not too close to the photometric limit of the 2MASS survey, the objects had to be brighter than $J = 16$. Elliptical objects may have inaccurate photometry and astrometry, hence selected objects had to have an axial ratio less than 1.4. Crowded regions near the galactic plane will produce many spurious detections of high proper motion objects. In order to reduce this crowding any object with $|b| < 15^\circ$ was excluded. The 2MASS Point Source Catalogue includes a parameter indicating the proximity of the nearest source; the Executive Summary for the catalogue states that any object closer than $6''$ to its nearest neighbouring source must be treated with caution. Hence such objects were excluded. Objects categorised by the 2MASS survey as being associated with extended sources or minor planets were also removed.

Objects which met these criteria were then subjected to a series of colour cuts.

In order to make the process as efficient as possible, only objects whose photometry suggested that they were M, L and T dwarfs were selected. In order to do this regions on a colour-colour diagram were marked out as likely to contain objects of a particular spectral type. Figure 1(a) shows these regions and Figure 1(b) shows the colours of objects taken from various surveys. Note that because there is an overlap between the mid M and mid T spectral classes on a colour-colour diagram any object that falls in this region is treated as if it could be either spectral type. Several early M dwarfs fall outside the bounds of the M dwarf region. When the full survey is produced this will have to be included in completeness estimates. Many T dwarfs fall outside the T dwarf region, this scatter is due to photometric errors. As we have set an I plate limit and an $I - J$ cut we will only identify the brightest T dwarfs. Dimmer objects have higher photometric errors and hence higher scatter. As these objects are already excluded due to our other

photometric cuts it does not matter if they fall outside the T dwarf region on the colour colour diagram.

Once objects have passed the colour cuts and been categorised it is then necessary to find if they are in fact moving with significant proper motions. Figure 2 shows the relative positional errors between SuperCOSMOS and 2MASS data. These were calculated by finding the positional shift of a set of 2MASS objects and calculating the mean error using a method of median absolute deviation. On the basis of these results it was decided to set one arcsecond (roughly $4\text{--}5\sigma$) as the lower “movement cut”. This ensures that only objects which appear to have moved from one epoch to the other are included in the sample. Any 2MASS object found to have an I plate counterpart within one arcsecond was discounted as not having a significant enough proper motion. This cut has implications for proper motion completeness which will have to be taken into account. Also to remove spurious detections due to bright stellar halos, any 2MASS object within $10''$ of an I plate image flagged as being close to a bright star was removed. Images blended on the I plates can cause positional offsets, hence any 2MASS object within $10''$ of a bright ($I < 14$) highly elliptical object or $6''$ of one de-blended by SuperCOSMOS software was removed.

2.2. Pairing with potential I plate counterparts

Once it is established that objects have appeared to move their I plate counterparts need to be identified. In order to do this a maximum proper motion of $\mu_{\text{max}} = 10''/\text{yr}$ was set meaning only I plate objects within a maximum radius $r_{\text{max}} = \mu_{\text{max}}\Delta t$ of the 2MASS object would be paired, where Δt is the time elapsed between the two observations. This maximum proper motion was chosen to be roughly equal to that of Barnard’s star, the star with the highest known proper motion, this allows us to identify very high proper motion stars without being swamped by spurious pairings. Once an I plate object was identified as within r_{max} of the target it had to pass a series of tests to prove it was indeed a plausible counterpart to the 2MASS object.

Firstly the I plate object had to have astrometry suggesting it was a high proper motion star. These data come from the calculated astrometric solutions in the SuperCOSMOS sky survey (Hambly et al. 2001). Any I plate object with an astrometric solution that places it at the same position (within the proper motion errors quoted in the astrometric solution) as the 2MASS object at the epoch of the 2MASS observation will be included as a possible counterpart. Many objects will either be so red that they will only appear on the I plates also objects moving with high proper motions will have moved too far to be paired in other bands so will be recorded as only being on the I plates. Hence any object appearing only on the I plates was included as a possible counterpart. Finally some I plate objects will be incorrectly paired with unre-




Fig. 1. The regions in the colour-colour diagram within which objects were deemed to be M, L or T dwarfs.

lated objects on the B_J or R plates and hence will have an inaccurate astrometric solution. Such objects will have a high value of χ^2_ν (χ^2 per degree of freedom), hence any object with $\chi^2_\nu > 2.0$ was included as a possible counterpart.

The I plate images also had to pass tests to indicate that they were good stellar images. Objects deblended by the SuperCOSMOS software were excluded as it is difficult to gather accurate astrometric information from them. Images with axial ratios higher than 1.7 were also excluded as they may have poor astrometry or may be blended objects which have not been separated by the software. I plate counterpart objects also had to be stellar in nature and not be in close proximity to bright stars.

Candidates for I plate counterpart images also had to pass a series of photometric tests. Firstly to ensure that the I plate counterparts were red enough to be true Low Mass Stars or Brown Dwarfs any object which had a UKST R plate magnitude had to pass the cut $(R-I) > 2$. As noted in Section 2.1, 2MASS objects were classified as being, M and L dwarfs, early T dwarfs, mid-late T Dwarfs or as lying in the overlap region on the colour-colour diagram be-




Fig. 2. The relative positional errors between SuperCOSMOS and 2MASS. The solid line represents the errors in Right Ascension while the dotted line shows those in Declination.

tween M dwarfs and mid-T Dwarfs. Objects classified as M and L dwarfs had to fulfill the condition $1.0 < I-J < 4.8$. Objects classified as early T Dwarfs or mid-late T dwarfs had to be within the range $4.0 < I-J < 5.5$. Any object falling in the overlap region between M dwarfs and mid-T dwarfs could be of either type so had to conform to the conservative cut of $1.0 < I-J < 5.5$.

In order to ensure that the I plate images selected were true counterparts and not unrelated objects two additional processes were carried out. Firstly it was checked that the I plate image was not associated with (within one arcsecond of) another 2MASS image. Secondly inspection by eye of the images ensured that only true high proper motion objects were selected.

3. Determination of Proper Motions

Objects in this paper are initially selected using proper motions calculated by finding the distance an object had moved between the two epochs and dividing by the time elapsed. Any object with such a calculated proper motion greater than $0.4''/\text{yr}$ was selected to have its proper

motion more accurately calculated. The SuperCOSMOS Sky Survey software calculates proper motions using B_J , I and two epoch R measurements for each object in the database. Hence, in theory, we should have access to astrometric solutions for all the objects selected so far. Unfortunately many do not have suitable solutions, stars with very high proper motions will have large shifts in their positions from plate to plate meaning they will often be too far apart from one epoch to the next for the the SSS software to identify them as the same object. Hence such stars will not have calculated proper motions. Some high proper motion objects may be spuriously paired with faint unrelated objects on one or more plates, producing an unreliable astrometric solution. Also the reddest objects may only appear on the I plate and hence will have no calculated astrometric solution. In order to make sure that all objects had a reliable astrometric solution a procedure to calculate the astrometric solution using only I plate and 2MASS data has been developed. Firstly a set of reference stars with I and J magnitudes similar to the star whose proper motion we wish to calculate was selected. These reference stars are then used to produce a linear fit between the plates, to account for plate to plate scaling and orientation errors. The stars positions are then corrected for plate to plate errors and used to find the proper motion. The plate to plate fit also provides an estimate of the typical error for objects near the target star and this is used to calculate the error in the proper motion. In the case where an object has a calculated astrometric solution both from the I plate-2MASS fitting procedure and from the SuperCOSMOS software the most accurate one is quoted. Unfortunately some objects did not have enough reference stars to produce a plate model. These had their proper motions found in a same way to those with enough reference stars with the only difference being the lack of corrections for plate to plate errors. The error estimate for these stars was calculated using the positional errors shown in Figure 2.

4. Results

Appendix A contains the astrometric data for all the objects identified with proper motions greater than $0.5''/\text{yr}$. There are 73 new objects and 71 objects identified by previous surveys. The I , J , H and K_s photometry for each object are also listed. Figure 3 shows the colours of the newly discovered objects. The reddest newly discovered object is SIPS0641–4322 which has colours suggesting it is an early L dwarf. Objects such as this would be difficult to find using purely optical proper motion surveys. Figure 4 shows histogram of the number of newly discovered and previously known objects vs. magnitude (J) and colour (I_J). It is clear that the newly discovered objects are generally fainter and redder than those previously known. Below we present notes on the most interesting individual newly identified objects identified.

Table 2. The full astrometric solution for SIPS1259-4336

Parameter	Fitted Value	Error	Units
RA on 01/01/2000	12 ^h 59 ^m 04 ^s .760	0.047 as	—
Dec on 01/01/2000	−43°36′24″.21	0.047 as	—
μ_α	1.105	0.004	as/yr
μ_δ	−0.262	0.004	as/yr
π	276	41	mas
χ^2_ν	0.65		

4.1. Notes On Individual Objects

4.1.1. SIPS1259–4336

Infrared colours indicate this object is an M8-9 dwarf 13. It is one magnitude fainter than the similar object SCR1845–6357 11 placing it at a distance of roughly 6pc. In addition its high proper motion suggests that it is a nearby star. Hence as there were several non-survey plates available for this object in the UKST plate library a trigonometric parallax measurement was carried out. This draws on techniques used in Deacon & Hambly (2001) and Deacon et al. (2005). Four of the non-survey plates were selected for their good parallax factors and were scanned on SuperCOSMOS and used along with the existing UKST R and ESO R plates. Details of the plates used are shown in Table 1.

The data from these plates were reduced using the same method as SCR1845-6357. A total of 148 reference stars were used to produce the plate to plate models. Once these had been applied to correct for plate to plate errors the astrometric solution could be found, this is shown in Table 2. The deviation from the proper motion is shown in Figure 6 with the parallax ellipse traced out by the object shown in Figure 7. To check for any systematic errors the reference stars were run through the fitting procedure. Their proper motions and parallaxes are shown in Figure 8 along with those for SIPS1259-4336. It is clear that there is no significant offset in the parallax and that SIPS1259-4336 is well separated from the mass of reference stars. The mean χ^2_ν for the reference stars was found to be 0.97 indicating good model fits. Spectroscopic data for this object were obtained from IRIS2 at the Anglo-Australian Telescope. These data were reduced and the spectrum in the J_l band is shown in Figure 5. Using the equivalent width of the potassium doublet as a diagnostic (Gorlova et al., 2003) we find that it is likely to be an M8 dwarf. The calculated parallax of SIPS1259-4336 of 276 ± 41 milliarcseconds relates to a distance of 3.62 ± 0.54 parsecs. This is nearer than expected as while the object is of similar colour to SCR1845-6357 it is one magnitude dimmer, hence it should be much further away. However if SCR1845-6357 is an unresolved double then the distance quoted here could be accurate.




Fig. 3. A two colour diagram for all the objects found in the survey. The star symbols represent new discoveries while the solid circles are previously known objects

4.1.2. SIPS1910–4132

When the candidate image for SIPS1910–4232 was inspected it was found that there were two objects moving with a similar proper motion. Astrometric solutions for these companion objects were found by the *I* plate-2MASS method as described in Section 3 and are shown in Table 3. It is clear that the three objects share a common proper motion and hence are likely to be a triple system. Table 4 shows the photometry for these objects. They all appear to be mid-late M dwarfs.

4.1.3. SIPS0251–0352

This object has the highest proper motion of the newly discovered objects. Its colours indicate that it is an early L dwarf (Kirkpatrick et al. 2000). Comparing its *J* magnitude with mean characteristics of L dwarfs in Vbra et al. (2004) shows that it appears to be within 20pc.

4.1.4. SIPS0052–6201

The astrometric solution for this object suggests that it has a common proper motion with LHS 128. LHS 128 is a K5 dwarf with a parallax of 52 mas (Perryman et al. 1997), if the two objects are part of the same system this yields a separation perpendicular to the line of sight of roughly 50,000 AU. It is also possible that the two objects are not part of a binary system but are simply members of the same moving group.

5. Completeness

5.1. Areal Completeness

As the final aim of this survey is to produce an estimate for the local space density of late M, L and T Dwarfs we need to analyse the area covered by our survey. We have already excluded objects near the Galactic Plane due to crowding. Our other main problem related to crowding is that of the proximity flag given in the 2MASS Point Source Catalogue. As advised by the 2MASS Executive Summary we have excluded all objects within 6 arcseconds of their nearest neighbour; this will present completeness problems in crowded regions. In order to produce an estimate as to the magnitude of these completeness problems we produced a series of simulations which allow us to calculate the probability of an object having a proximity flag greater than 6 arcseconds for a given sky surface density of 2MASS images. To do this a number of images were randomly placed in a one arcminute square box. If any of these fell within 6 arcseconds of the centre the test was deemed to have failed. However if no image fell within 6 arcseconds of the centre the test was deemed to have succeeded. The probability of detection is thus the number of successful tests divided by the total number of tests. Simulations were carried out for different numbers of stars per square arc minute ranging from 1 to 100. The results of these simulations are shown in Figure 9. In order to calculate the probability of objects being identified in particular regions of the sky, the number of 2MASS images per square arcminute and hence the probability of detection




Fig. 5. The normalized spectrum of SIPS1259–4336 (top) along with that of an A0 standard star. The potassium doublet at 1.25 microns can clearly be seen.

were calculated for each UKST field.

In addition there will be areal completeness problems caused by areas of UKST plates which are near to bright stars. Luckily the size of these areas is calculated in the post processing of each SuperCOSMOS scan. Hence we simply multiply this probability of detection that for the 2MASS images to yield an estimate of the areal completeness for each field. Figure 10 shows these data as a grey-scale plot. It is clear that the galactic plane is crowded and hence unsuitable for inclusion in this survey.

Further completeness issues will need to be addressed before a proper estimate of the the luminosity function (and hence mass function) of the objects included here can be determined. Proper motion completeness will need to be simulated for, with the main factors being the one arcsecond “movement” cut and the lower proper motion limit. Also photometric completeness must be estimated taking both the 2MASS colour cut and the limit on the *I* plate magnitude.




Fig. 6. The deviation from the proper motion in RA and Dec plotted against time. The line shown is the path of parallax motion predicted by the astrometric solution

5.2. Comparison with other studies

In order to make clear the rough extent of the currently unquantified incompleteness due to astrometric and deblending cuts the survey was compared to the Luyten Half-Arcsecond Catalogue (Luyten 1979). Using the accurate positions of Bakos, Sahu and Nemeth (2002) the 2MASS data for the LHS objects was extracted. These objects were then subjected to the photometric, crowding and bright star cuts as the survey data. Figure 11 shows the number that could be detected and those that were. It is clear that the survey is somewhat incomplete, with only 43 of the 200 Luyten objects recovered. This is mainly due to many 2MASS and *I* observations being taken at virtually the same time leading to a short proper motion baseline but deblended and elliptical objects will also play a role. However this is not an accurate estimate of the incompleteness of the survey. As the Luyten survey will be unquantifiably incomplete itself it will be necessary to produce simulations for both the photometric and astrometric incompleteness in order to gain a complete picture of the total incompleteness of the survey.




Fig. 7. The parallax ellipse traced out by SIPS1259-4336; the dotted lines represent one sigma upper and lower limits on the parallax.




Fig. 8. Comparison of the fitted parallax and proper motions for the reference stars with SIPS1259-4336.

6. Conclusion and Further Work

We have utilised wide field surveys to discover 72 new objects with proper motions greater than $0.5''/\text{yr}$. Of these objects we have discovered that one of them has a significant trigonometric parallax of $\pi = 276 \pm 41$ milliarcseconds yielding a distance of $3.62 \pm 0.54 \text{ pc}$.

In this paper our lower proper motion sets a bias towards nearer and hence brighter objects. In future papers we hope to detect objects with much lower proper motions allowing us to expand our search volume greatly. We will also attempt to identify ultracool members of different kinematic populations by means of reduced proper mo-




Fig. 9. The probability of detection plotted against the number of 2MASS images per square arc minute.

tions. Once the SuperCOSMOS Sky Survey is completed on the northern sky we hope to make use of these data too. Finally we hope to utilise the data yielded to these studies along with more substantial magnitude, proper motion and volume completeness estimates to calculate the local space density of late M,L and T dwarfs.

Acknowledgements. The authors would like to thank Sue Tritton and Mike Read for their help in selecting plates, Harvey MacGillivray and Eve Thomson for their prompt scanning of the material on SuperCOSMOS, David Bacon for his useful discussions on statistics, Mairi Brookes and Jessica Skelton for their discussion on spectroscopy and to Stuart Ryder for taking spectroscopic observations. This publication makes use of data products from the Two Micron All Sky Survey, which is a joint project of the University of Massachusetts and the Infrared Processing and Analysis Center/California Institute of Technology, funded by the National Aeronautics and Space Administration and the National Science Foundation. SuperCOSMOS is funded by a grant from the UK Particle Physics and Astronomy Research Council.

References

- Bakos, G., Sahu, K.C., Nemeth, P., AJSS, 141, 1, 187, 2002
- Burgasser, A.J., McElwain, M.W., Kirkpatrick, J.D., Cruz, K.L., Tinney, C.G., Reid, I.N., AJ, 127, 5, 2856, 2004

- Burgasser, A.J. et al., *ApJ*, 564, , 421, 2002
- Dahn, C.C. et al., *AJ*, 124, 2, 1170, 2002
- Deacon, N.R, Hambly, N.C., *A&A*, 380, 148, 2001
- Deacon, N.R, Hambly, N.C., Henry, T.J., Subasavage, J., Brown, M., Jao, W.C., *AJ* accepted, astro-ph/0409582, 2005
- Geballe, T.R. et al. *ApJ*, 564, 1, 466, 2002
- Gorlova, N. I., Meyer, M. R., Rieke, G. H., Liebert, J., *ApJ*, 593, 2, 1074, 2003
- Hambly, N.C., MacGillivray, H.T., Read, M.A., Tritton, S.B., Thomson, E.B., Kelly, B.D., Morgan, D.H., Smith, R.E., Driver, S.P., Williamson, J., Parker, Q.A., Hawkins, M.R.S., Williams, P.M., Lawrence, A., *MNRAS*, 326, 4, 1279, 2001
- Hambly, N. C.; Davenhall, A. C.; Irwin, M. J.; MacGillivray, H. T., *MNRAS*, 326, 4, 1315, 2001
- Hambly, N.C., Henry, T.J., Subasavage, J., Brown, M., Jao, W.C., *AJ*, 128:437-447, 2004
- Kendall, T. R., Delfosse, X., Martin, E. L., Forveille, T., *A&A* 416, L17, 2004
- Kirkpatrick, J.D., Reid, I.N., Liebert, J., Gizis, J.E., Burgasser, A.J., Monet, D.G., Dahn, C., Nelson, B., Williams, R.J., *AJ*, 120, 1, 447, 2000
- Lepine, S., Shara, M.M., *AJ*, 124, 2002
- Luyten Half Arcsecond Catalogue, Luyten, W.J., University of Minnesota, Minneapolis, 1979
- Perryman, M. A. C., Lindegren, L., Kovalevsky, J., Hoeg, E., Bastian, U., Bernacca, P. L., Crz, M., Donati, F., Grenon, M., van Leeuwen, F., van der Marel, H., Mignard, F., Murray, C. A., Le Poole, R. S., Schrijver, H., Turon, C., Arenou, F., Froeschl, M., Petersen, C. S., *A&A*, 323, L49, 1997
- Pokorny, R.S., Jones, H.R.A., Hambly, N.C., *A&A*, 397, 575-584, 2003
- Ruiz, M.T., Wischnjewsky, M., Rojo, P.M., Gonzalez, L.E., *ApJSS*, 133, 119-160, 2001
- Subasavage, J.P., Henry, T.J., Hambly, N.C., Brown, M.A., Jao, W.C., *AJ* submitted, 2005
- Scholz, R.D., Ibata, R., Irwin, M., Lehmann, I., Salvato, M., Schweitzer, A., *MNRAS*, 329, 109-114, 2002
- Scholz, R.D., McCaughrean, M.J., Lodieu, N., Kuhlbrodt, B., *A&A*, v. 398, L29-L33, 2004
- Seifahrt, A., Neuhuser, R., Mugrauer, M., *A & A*, 421, 255, 2004
- Teegarden, B.J., Pravdo, S.H., Hicks, M., Lawrence, K., Shaklan, S.B., Covey, K., Fraser, O., Hawley, S.L., McGlynn, T., Reid, I.N., *ApJ*, 589, L51-L53, 2003
- Vbra, F.J. et al., *AJ*, 127, 5, 2948, 2004
- Wallace, P.T., Starlink User Note No. 67.42: SLALIB: Positional Astronomy Library, CCLRC/Rutherford Appleton Laboratory, PPARC, 1998
- Wroblewski, H., Costa, E., *A&A*, 367, 725-728, 2001
- The 2MASS Point Source Catalogue Executive Summary, <http://pegasus.phast.umass.edu/>




Fig. 4. Histograms showing the magnitude (J) and colour ($I - J$) of both the newly discovered and previously

Table 1. Schmidt photographs used in this study; relative astrometric quality is indicated (see text). One plate was excluded due to poor astrometric quality.

Plate No.	Date (dd/mm/yy)	LST	Zenith Angle	Emulsion	Filter	Exp. (min)	σ_x (mas)	σ_y (mas)	Material
Plates Used									
R 5108	25/06/79	1254	12.5°	IIIaF	RG 630	800	64	65	Original non-survey plate
ESOR4809	19/05/82	1200	18.6°	IIIaF	RG630	120	63	49	Copy of ESO survey plate
OR14770	12/02/92	1245	12.6°	IIIaF	OG 590	600	41	49	Original survey plate
OR16928	26/01/96	1142	19.6°	4415	OG 590	150	68	67	Original non-survey film
OR17069	26/04/96	1322	13.1°	IIIaF	OG 590	100	56	55	Original non-survey plate
OR17997	28/04/98	1314	12.7°	IIIaF	OG 590	600	77	65	Original non-survey plate

Table 3.

Name	Position (J2000)	Epoch	μ_{tot}	PA	σ_μ
SIPS1910–4133A	19 10 34.50 -41 33 32.5	1983.346	0.742	174.4	0.017
SIPS1910–4133B	19 10 45.87 -41 33 28.9	1983.346	0.748	173.0	0.021
SIPS1910–4132C	19 10 33.49 -41 32 38.8	1983.346	0.738	174.7	0.016

Table 4.

Object	I	J	H	K_S
SIPS1910–4133A	11.008	9.851	9.245	9.032
SIPS1910–4133B	11.738	10.61	10.002	9.739
SIPS1910–4132C	12.577	11.147	10.552	10.249




Fig. 10. The areal completeness of the survey. The scale shows the probability that an area is free from both crowding and bright stars. The solid lines marked represent the galactic latitude cut.




Fig. 11. The number of Luyten objects that could possibly be detected (grey) and those that were (solid).

Appendix A: Catalogue tables

Table A.1 contains details of objects previously undiscovered with proper motions greater than $0.5''/\text{yr}$. Objects with proper motions taken from the SuperCOSMOS sky survey are marked with “a”, while objects with proper motions calculated from 2MASS and SuperCOSMOS *I* data are marked with “b”. Objects marked “c” lacked sufficient reference stars to produce a plate to plate fit. For objects with LHS numbers see Luyten (1979) and references therein. Objects with a CE prefix can be found in Ruiz et al. (2001), objects with LEHPM numbers are in Pokorny, Jones & Hambly (2003) and APMPM objects are to be found in Scholz et al. (2002). For objects with a quoted 2MASS designation see Burgasser et al. (2004) and references therein. For objects with a quoted DENIS designation see Kendall et al. (2004) and references therein. Finally for objects marked WT see Wroblewski & Costa (2001) and references therein.

Table A.1. Objects marked * were simultaneously discovered by Subasavage et al. (2005).

Name	Position (J2000)	Epoch	μ_{tot}	PA (arcseconds/yr)	σ_μ (Degrees)	Proper Motion Source
SIPS0308–8212 *	03 08 53.50 -82 12 34.9	1990.818	0.511	28.9	0.017	b
SIPS1240–8209 *	12 40 53.21 -82 09 03.9	1991.100	0.541	276.7	0.019	b
SIPS2130–7710 *	21 30 06.31 -77 10 36.2	1995.669	0.653	118.1	0.037	b
SIPS1633–7603	16 33 57.58 -76 03 54.4	1996.397	0.510	198.9	0.055	b
SIPS2150–7520	21 50 14.20 -75 20 34.7	1993.733	1.019	106.0	0.050	b
SIPS0452–7322 *	04 52 04.61 -73 22 03.6	1978.930	0.582	54.2	0.015	b
SIPS0321–7046	03 21 14.80 -70 46 11.5	1990.979	0.553	40.3	0.025	b
SIPS1924–6920	19 24 35.60 -69 20 13.0	1996.357	0.817	148.3	0.042	b
SIPS2032–6918	20 32 32.62 -69 18 56.4	1994.650	0.585	147.2	0.027	b
SIPS1932–6506	19 32 36.39 -65 06 44.9	1994.353	0.510	168.3	0.030	b
SIPS0052–6201	00 52 14.49 -62 01 55.1	1994.621	1.123	82.8	0.050	b
SIPS1943–6125	19 43 33.11 -61 25 37.0	1993.637	0.791	179.0	0.034	b
SIPS0523–5608	05 23 03.82 -56 08 42.5	1992.033	0.683	341.8	0.024	b
SIPS1936–5502	19 36 01.69 -55 02 30.7	1996.559	0.836	133.8	0.051	b
SIPS2053–5409	20 53 03.74 -54 09 33.7	1994.688	0.722	158.1	0.033	b
SIPS2242–4514	22 42 04.80 -45 14 57.1	2000.731	0.836	280.5	0.225	b
SIPS1410–4425	14 10 41.08 -44 25 55.9	1982.437	0.537	195.9	0.024	b
SIPS1259–4336	12 59 03.87 -43 36 22.1	1991.226	1.133	104.0	0.021	b
SIPS0641–4322	06 41 18.30 -43 22 36.2	1994.212	0.689	19.2	0.040	b
SIPS1337–4311	13 37 56.25 -43 11 28.2	1994.453	0.552	227.4	0.046	b

Table A.1.

Name	Position (J2000)	Epoch	μ_{tot}	PA (arcseconds/yr)	σ_μ (Degrees)	Proper Motion Source
SIPS1910–4132	19 10 33.49 -41 32 38.8	1983.346	0.738	174.7	0.016	b
SIPS1231–4018	12 31 21.55 -40 18 37.1	1993.249	0.679	274.0	0.033	b
SIPS1149–4012	11 49 50.12 -40 12 42.0	1991.223	0.566	253.2	0.034	b
SIPS0013–3921	00 13 00.54 -39 21 19.9	1994.648	0.522	102.7	0.033	b
SIPS1251–3846	12 51 32.08 -38 46 13.1	1994.297	0.593	278.9	0.036	b
SIPS0031–3840	00 31 18.84 -38 40 35.1	1990.817	0.568	96.8	0.035	c
SIPS1039–3819	10 39 32.57 -38 19 57.2	1992.080	0.724	203.6	0.035	b
SIPS1338–3752	13 38 28.14 -37 52 50.2	1984.304	1.278	269.6	0.014	b
SIPS1141–3624	11 41 21.05 -36 24 38.3	1988.154	0.556	57.1	0.013	b
SIPS1541–3609	15 41 19.25 -36 09 10.5	1994.407	0.507	263.7	0.038	b
SIPS1342–3534	13 42 22.47 -35 34 47.9	1984.304	0.931	259.1	0.015	b
SIPS2050–3358	20 50 44.01 -33 58 37.5	1986.563	0.510	120.4	0.022	b
SIPS0417–3211	04 17 58.82 -32 11 49.4	2000.967	0.855	283.7	0.111	b
SIPS2346–3153	23 46 54.50 -31 53 50.5	1992.551	0.649	134.9	0.043	b
SIPS0405–3138	04 05 29.44 -31 38 44.6	2000.967	0.846	261.2	0.053	b
SIPS2314–2929	23 14 26.40 -29 29 52.8	1994.569	0.620	132.7	0.043	a
SIPS1529–2907	15 29 14.08 -29 07 28.8	1991.452	1.078	185.8	0.025	b
SIPS1548–2859	15 48 21.55 -28 59 33.9	1994.349	0.546	219.0	0.052	b
SIPS2313–2826	23 13 59.28 -28 26 36.9	1994.653	0.825	117.5	0.054	b
SIPS0422–2802	04 22 31.25 -28 02 30.6	2000.967	0.914	281.5	0.077	b
SIPS2231–2756	22 31 22.38 -27 56 47.3	1999.691	1.677	105.8	0.407	b
SIPS1025–2730	10 25 55.01 -27 30 59.6	1996.227	0.502	223.9	0.041	b
SIPS2157–2726	21 57 54.53 -27 26 28.8	1994.692	0.855	131.4	0.053	b
SIPS2308–2721	23 08 11.23 -27 21 59.1	1994.771	0.597	124.7	0.067	b
SIPS1019–2707	10 19 24.68 -27 07 17.3	1994.371	0.582	272.3	0.044	b
SIPS1540–2613	15 40 30.18 -26 13 35.4	1993.328	1.623	224.9	0.021	b
SIPS1424–2535	14 24 53.94 -25 35 17.9	1997.238	0.641	105.5	0.175	b
SIPS1627–2518	16 27 49.75 -25 18 13.5	1982.568	0.582	210.8	0.015	b
SIPS1513–2243	15 13 59.26 -22 43 47.6	1992.242	0.574	206.4	0.026	b
SIPS1250–2121	12 50 52.43 -21 21 11.4	1993.399	0.578	126.0	0.042	c
SIPS0921–2104	09 21 13.97 -21 04 38.1	1993.169	0.965	162.6	0.016	b
SIPS0004–2058	00 04 41.17 -20 58 30.3	1994.733	0.826	90.6	0.107	a
SIPS2359–2007	23 59 57.48 -20 07 37.6	1994.733	0.798	127.7	0.040	b
SIPS2210–1952	22 10 50.04 -19 52 22.1	1994.771	0.830	188.6	0.051	b
SIPS1048–1925	10 48 18.98 -19 25 33.3	1996.388	0.957	264.4	0.112	b
SIPS1121–1653	11 21 09.57 -16 53 52.1	1993.396	0.602	204.8	0.0473	b
SIPS0227–1624	02 27 10.17 -16 24 46.2	1994.927	0.592	120.8	0.019	b
SIPS1109–1606	11 09 27.58 -16 06 50.4	1994.139	0.520	233.9	0.050	b
SIPS0933–1602	09 33 49.78 -16 02 50.7	1997.000	1.357	188.3	0.072	b
SIPS0050–1538	00 50 24.35 -15 38 20.0	2001.796	0.674	227.3	0.042	b
SIPS0258–1220	02 58 35.03 -12 20 18.3	2001.805	0.827	91.4	0.042	b
SIPS1413–1201	14 13 05.33 -12 01 22.4	1988.545	0.715	237.2	0.025	b
SIPS1542–1007	15 42 25.44 -10 07 02.3	1996.302	0.552	166.0	0.062	b
SIPS2025–0835	20 25 47.58 -08 35 29.0	1994.661	0.578	151.8	0.052	b
SIPS0320–0446	03 20 28.43 -04 46 33.4	1994.698	0.678	190.6	0.038	b
SIPS1956–0422	19 56 50.17 -04 22 39.3	1993.634	0.563	115.0	0.026	b
SIPS0820–0355	08 20 44.43 -03 55 04.5	1996.224	0.594	146.6	0.123	b
SIPS0251–0352	02 51 14.53 -03 52 36.3	1993.690	2.185	149.3	0.057	b
SIPS0346–0218	03 46 11.89 -02 18 18.1	1986.891	0.850	155.4	0.021	b
SIPS0937–0214	09 37 31.35 -02 14 32.4	1987.099	0.528	127.6	0.025	b
SIPS2343–0155	23 43 04.40 -01 55 06.0	1999.786	1.359	279.4	0.288	b
SIPS1512–0112	15 12 17.92 -01 12 28.3	1990.489	0.671	191.6	0.031	b

Table A.2.

Name	Position (J2000)	Epoch	μ_{tot}	PA (arcseconds/yr)	σ_μ (Degrees)	Proper Motion Source
SCR 1845–6357	18 45 02.32 -63 57 52.6	1992.586	2.541	74.1	0.153	a
LHS 532	22 56 25.04 -60 03 44.4	1994.648	1.020	206.6	0.024	b
LHS 3492	19 51 31.35 -50 55 33.1	1994.366	0.815	192.2	0.033	b
APMPM J04052–5819	04 51 37.56 -58 18 44.5	1989.038	0.701	192.6	0.025	b
WT 60	01 52 00.00 -57 47 55.7	1994.858	0.636	216.5	0.037	b
ϵ Indi B	22 04 09.46 -56 46 52.6	1997.771	4.840	120.8	0.048	c
APMPM J0524–5607	05 23 30.59 -56 07 06.4	1992.033	0.782	5.8	0.029	b
LEHPM 1683	01 34 59.74 -55 11 34.9	1994.856	0.644	135.0	0.042	b
APMPM J 0247–5257	02 47 20.63 -52 56 45.9	1997.723	0.776	47.1	0.102	a
LHS 1471	02 55 14.27 -51 40 21.8	1996.964	0.656	70.5	0.094	a
LEHPM 3396	03 34 08.52 -49 53 39.5	1984.915	2.448	77.6	0.055	a
DENIS J0255–4700	02 55 03.18 -47 00 48.6	1994.809	1.197	120.5	0.126	a
LHS 3971	23 34 15.53 -40 44 03.8	1984.625	0.857	105.2	0.023	a
LHS 1502	03 08 49.00 -40 06 40.4	1995.784	0.739	142.9	0.058	b
DENIS J1048–3956	10 48 15.39 -39 55 59.3	1992.080	1.541	231.0	0.013	b
LHS 1490	03 02 06.56 -39 50 49.3	1995.784	0.792	221.5	0.025	b
LHS 539	23 15 51.15 -37 33 32.1	1995.625	1.352	71.1	0.084	b
2MASSW J1155395–372735	11 55 39.51 -37 27 32.5	1996.189	0.868	172.5	0.039	b
LHS 1531	03 19 17.03 -37 03 42.0	1995.989	0.818	124.8	0.039	b
LEHPM 3070	03 06 11.65 -36 47 50.3	1995.989	0.571	179.8	0.243	a
CE 136	10 11 13.49 -35 36 21.6	1993.325	0.509	293.2	0.030	b
LHS 1192	01 05 52.14 -34 34 48.5	1991.582	0.607	92.2	0.035	b
LHS 1355	02 12 39.83 -33 52 06.1	2000.723	1.045	93.5	0.083	b
LHS 2506	12 03 58.50 -33 01 26.8	1996.189	0.768	275.4	0.033	b
CE 220	10 45 34.15 -32 49 53.5	1993.246	0.551	306.6	0.046	b
APMPM J1407–3018	14 06 49.79 -30 18 27.6	1993.208	0.852	264.3	0.028	b
LHS 3566	20 39 23.68 -29 26 30.6	1994.538	0.832	154.1	0.041	b
LHS 3662	21 23 07.74 -28 09 54.4	1994.773	0.715	132.9	0.057	b
APMPM J1224–2758	12 23 56.82 -27 57 48.4	1993.243	1.247	282.9	0.027	b
APMPM J2331–2750	23 31 21.70 -27 49 53.6	1994.569	0.781	1.439	0.136	a
LHS 2487	11 58 57.02 -27 43 07.5	1993.246	0.501	295.4	0.031	b
DENIS J1456–2747	14 56 01.44 -27 47 34.7	1997.361	0.821	197.3	0.028	b
LHS 2597	12 39 36.56 -26 58 10.6	1997.361	0.713	259.5	0.036	b
LHS 2928	14 30 18.06 -24 03 15.3	1997.238	0.637	228.8	0.062	a
LEHPM 162	00 05 48.18 -21 57 19.0	1994.733	0.778	104.7	0.045	b
LHS 3970	23 33 40.41 -21 33 51.9	1995.628	0.794	103.2	0.027	b
LHS 1069	00 24 39.49 -20 54 09.1	1995.669	0.507	198.7	0.042	b
LHS 105	00 09 16.95 -19 42 34.5	1994.733	1.093	60.8	0.036	b
LHS 3718	21 49 23.86 -18 10 36.6	1994.618	0.590	206.4	0.052	b
LHS 3799	22 23 06.98 -17 36 26.0	1999.606	0.929	164.5	0.240	c
LHS 138	01 12 30.28 -16 59 59.1	1995.822	1.386	63.6	0.160	a
2MASS J15074769–1627386	15 07 47.70 -16 27 37.5	1997.244	1.038	182.4	0.255	c
LHS 1396	02 23 01.78 -16 16 43.5	1994.927	0.781	125.7	0.033	b
LHS 2397	11 20 26.46 -14 39 59.8	1993.396	0.509	215.1	0.038	b
LHS 1462	02 52 39.40 -14 31 44.6	2001.805	0.865	195.7	0.042	a
LHS 500	20 55 35.92 -14 03 49.2	1987.734	1.484	107.8	0.013	b
LHS 523	22 28 54.36 -13 25 19.1	1999.601	1.222	209.1	0.129	b
LHS 3056	15 19 11.90 -12 45 05.6	1996.394	0.772	236.6	0.077	b
LHS 3935	23 21 13.24 -11 37 23.1	1994.776	0.760	104.3	0.048	b
LHS 3911	23 13 19.35 -11 06 17.1	1994.776	0.926	101.9	0.072	b
2MASSW J1555157–095605	15 55 15.28 -09 55 59.8	1992.310	1.273	131.1	0.022	b

Table A.2.

Name	Position (J2000)	Epoch	μ_{tot}	PA (arcseconds/yr)	σ_μ (Degrees)	Proper Motion Source
LHS 3735	21 57 39.27 -09 28 13.6	2000.723	0.541	248.9	0.061	b
LHS 353	13 30 03.00 -08 42 24.3	1997.320	1.362	249.2	0.041	b
LHS 176	03 35 38.61 -08 29 22.9	1999.995	1.748	112.2	0.050	b
LHS 424	16 55 35.58 -08 23 35.1	1993.632	1.163	221.503	0.031	b
LHS 2	00 06 43.16 -07 32 18.5	2000.723	2.063	201.3	0.120	c
LHS 5217	12 31 23.99 -06 38 00.2	1997.238	0.533	117.0	0.045	b
LHS 3149	16 04 20.02 -06 16 39.6	1992.504	0.867	182.9	0.028	b
DENIS J1539–0520	15 39 41.61 -05 20 43.3	1992.506	0.644	83.3	0.042	c
LHS 3611	21 01 53.75 -05 23 40.1	1995.510	0.528	179.6	0.025	a
LHS 1785	05 47 08.98 -05 12 09.3	1996.068	0.787	136.4	0.037	a
LHS 2419	11 31 26.85 -05 03 36.3	1995.138	0.582	278.7	0.029	a
2MASS J23062928–0502285	23 06 28.98 -05 02 26.0	1993.730	1.042	119.7	0.025	b
LHS 17	02 46 14.21 -04 59 08.8	1993.690	2.488	137.9	0.041	b
DENIS J0012–0457	00 13 46.38 -04 57 36.2	1994.651	0.685	107.4	0.039	b
LHS 1363	02 14 12.38 -03 57 42.8	1994.659	0.504	105.0	0.079	b
LHS 2641	12 49 48.72 -03 17 32.3	1997.243	0.608	195.6	0.049	a
LHS 3104	15 42 30.00 -03 15 51.7	1992.506	0.504	253.7	0.027	b
LHS 1332	02 04 27.97 -01 52 49.9	1987.822	0.799	225.2	0.013	b
LHS 1415	02 50 06.29 -01 50 32.8	1986.932	0.801	76.9	0.021	b
LHS 5165	10 03 19.26 -01 05 08.1	1997.123	0.538	284.0	0.060	b
LSR J1610–0040	16 10 28.93 -00 40 54.3	2000.242	1.719	226.6	0.088	b

Table A.3.

Object	I	J	H	K_S
SIPS0308–8212	13.087	11.704	11.145	10.894
SIPS1240–8209	12.490	10.855	10.196	9.933
SIPS2130–7710	13.476	11.292	10.666	10.365
SIPS1633–7603	16.655	14.261	13.644	13.298
SIPS2150–7520	17.260	14.056	13.176	12.673
SIPS0452–7322	13.621	11.976	11.442	11.124
SIPS0321–7046	11.746	10.394	9.840	9.574
SIPS1924–6920	14.267	12.392	11.908	11.607
SIPS2032–6918	16.298	13.636	12.983	12.579
SIPS1932–6506	14.784	12.664	12.048	11.719
SIPS0052–6201	14.184	12.153	11.742	11.370
SIPS1943–6125	16.115	13.199	12.700	12.287
SIPS0523–5608	15.580	13.554	12.948	12.602
SIPS1936–5502	17.474	14.486	13.628	13.046
SIPS2053–5409	13.916	12.170	11.656	11.345
SIPS2242–4514	17.501	15.981	15.586	15.120
SIPS1410–4425	17.980	15.874	15.066	14.544
SIPS1259–4336	12.799	10.534	9.954	9.520
SIPS0641–4322	17.170	13.751	12.941	12.451
SIPS1337–4311	12.850	11.701	11.112	10.794
SIPS1910–4132	12.577	11.147	10.552	10.249
SIPS1231–4018	11.538	10.444	9.940	9.635
SIPS1149–4012	17.269	13.999	13.322	12.834
SIPS0013–3921	17.551	14.787	14.216	13.809
SIPS1251–3846	13.758	12.086	11.558	11.246
SIPS0031–3840	17.442	14.101	13.399	12.924
SIPS1039–3819	12.535	10.846	10.301	9.998
SIPS1338–3752	13.141	11.747	11.281	10.965
SIPS1141–3624	10.108	8.490	7.967	7.699
SIPS1541–3609	14.467	11.970	11.450	11.109
SIPS1342–3534	14.790	13.755	13.216	12.935
SIPS2050–3358	15.858	13.815	13.146	12.854
SIPS0417–3211	17.562	14.864	14.196	13.739
SIPS2346–3153	15.838	13.279	12.680	12.198
SIPS0405–3138	13.978	12.924	12.282	11.989
SIPS2314–2929	17.329	14.608	14.101	13.682
SIPS1529–2907	15.475	13.319	12.855	12.499
SIPS1548–2859	15.468	13.118	12.551	12.260
SIPS2313–2826	14.499	12.660	12.149	11.878
SIPS0422–2802	16.250	14.755	14.091	13.775
SIPS2231–2756	17.859	15.808	15.318	14.903
SIPS1025–2730	16.699	14.551	14.048	13.625
SIPS2157–2726	17.914	14.672	13.887	13.401
SIPS2308–2721	17.819	14.662	13.833	13.332
SIPS1019–2707	16.753	13.526	12.906	12.471
SIPS1540–2613	14.600	11.646	11.145	10.730
SIPS1424–2535	17.751	15.665	15.123	14.784
SIPS1627–2518	17.706	14.888	14.368	14.016
SIPS1513–2243	17.064	14.196	13.601	13.231
SIPS1250–2121	13.336	11.160	10.550	10.128
SIPS0921–2104	16.031	12.779	12.152	11.690
SIPS0004–2058	15.647	12.404	11.834	11.396
SIPS2359–2007	17.380	14.382	13.623	13.248
SIPS2210–1952	16.456	14.000	13.498	13.151
SIPS1048–1925	17.830	14.876	14.276	13.692
SIPS1121–1653	13.856	12.553	12.073	11.762
SIPS0227–1624	16.711	13.573	12.630	12.143

Table A.4.

Object	<i>I</i>	<i>J</i>	<i>H</i>	<i>K_S</i>
SIPS1109–1606	17.856	14.970	14.348	13.892
SIPS0933–1602	15.307	12.720	12.232	11.890
SIPS0050–1538	16.880	13.779	13.077	12.647
SIPS0258–1220	15.822	13.921	13.410	13.030
SIPS1413–1201	10.446	9.040	8.453	8.163
SIPS1542–1007	16.859	14.372	13.801	13.437
SIPS2025–0835	15.699	13.153	12.403	12.029
SIPS0320–0446	16.558	13.259	12.535	12.134
SIPS1956–0422	16.949	13.441	12.921	12.510
SIPS0820–0355	17.319	15.884	15.257	15.025
SIPS0251–0352	16.214	13.059	12.254	11.662
SIPS0346–0218	15.255	13.130	12.642	12.346
SIPS0937–0214	16.979	14.410	13.883	13.585
SIPS2343–0155	17.868	15.598	15.044	14.757
SIPS1512–0112	16.607	15.244	14.873	14.469

Table A.5.

Object	<i>I</i>	<i>J</i>	<i>H</i>	<i>K_S</i>
SCR1845–6357	12.528	9.544	8.967	8.508
LHS 532	10.653	8.984	8.360	8.108
LHS 3492	12.288	10.970	10.493	10.188
APMPM J0452–5819	13.980	11.691	11.089	10.705
WT 60	13.213	11.450	10.846	10.568
ϵ Indi B	16.567	11.908	11.306	11.208
APMPM J0524–5607	13.960	12.337	11.715	11.370
LEHPM 1683	15.211	13.227	12.709	12.427
APMPM J0247–5257	13.097	11.748	11.184	10.873
LHS 1471	11.125	9.938	9.366	9.082
LEHPM 3396	13.916	11.376	10.823	10.392
DENIS 0255–4700	17.025	13.246	12.204	11.558
LHS 3971	12.683	11.248	10.713	10.435
LHS 1502	13.925	12.623	12.105	11.840
DENIS 1048–3956	12.097	9.538	8.905	8.447
LHS 1490	11.926	10.705	10.178	9.885
LHS 539	11.556	10.403	9.872	9.592
2MASSW J1155395–372735	16.370	12.811	12.040	11.462
LHS 1531	13.367	12.160	11.676	11.344
LEHPM 3070	13.764	11.690	11.068	10.631
CE 136	12.750	11.666	11.070	10.807
LHS 1192	12.808	11.160	10.569	10.272
LHS 1355	12.668	11.141	10.625	10.319
LHS 2506	10.618	9.529	9.073	8.744
CE 220	11.055	9.9869	9.233	8.966
APMPM J1407–3018	13.538	11.361	10.687	10.366
LHS 3566	13.682	11.357	10.743	10.367
LHS 3662	12.589	11.062	10.523	10.247
APMPM J1224–2758	14.151	11.977	11.401	11.069
APMPM J2331–2750	14.404	11.646	11.055	10.651
LHS 2487	14.359	12.763	12.238	11.963
DENIS 1456–2747	16.503	13.250	12.655	12.189
LHS 2597	11.424	10.042	9.581	9.233
LHS 2928	11.857	10.742	10.237	9.917
LEHPM 162	16.461	13.274	12.617	12.201
LHS 3970	13.725	11.858	11.312	10.929
LHS 1069	14.627	12.873	12.312	12.014
LHS 105	12.532	10.883	10.327	10.074
LHS 3718	13.994	12.532	12.052	11.747
LHS 3799	9.268	8.242	7.638	7.319
LHS 138	8.523	7.258	6.749	6.420
2MASS J15074769–1627386	16.134	12.830	11.895	11.312
LHS 1396	15.159	13.490	13.027	12.645
LHS 2397	14.877	13.444	12.903	12.641
LHS 1462	14.094	12.695	12.227	11.900
LHS 500	11.248	9.717	9.218	8.915
LHS 523	12.639	10.768	10.217	9.843
LHS 3056	9.640	8.507	7.862	7.582
LHS 3935	13.986	12.296	11.761	11.512
LHS 3911	12.866	11.125	10.607	10.299
2MASSW J1555157–095605	16.057	12.557	11.984	11.443

Table A.6.

Object	<i>I</i>	<i>J</i>	<i>H</i>	<i>K_S</i>
LHS 3735	12.019	10.450	9.822	9.537
LHS 353	10.731	9.599	9.047	8.749
LHS 176	11.596	10.377	9.801	9.456
LHS 424	11.595	9.776	9.201	8.816
LHS 2	9.876	8.323	7.792	7.439
LHS 5217	12.527	11.328	10.747	10.434
LHS 3149	11.683	10.452	9.880	9.548
DENIS 1539–0520	17.805	13.922	13.060	12.575
LHS 3611	14.00	12.839	12.33	12.006
LHS 1785	11.096	10.039	9.514	9.177
LHS 2419	15.216	13.682	13.171	12.860
2MASS J23062928–0502285	13.580	11.354	10.718	10.296
DENIS 0012–0457	13.925	11.462	10.866	10.479
LHS 17	12.102	10.970	10.499	10.152
LHS 1363	11.956	10.481	9.858	9.485
LHS 2641	11.874	10.759	10.261	9.925
LHS 3104	14.941	12.657	12.162	11.810
LHS 1332	10.744	9.585	9.092	8.804
LHS 1415	14.520	12.914	12.430	12.119
LHS 5165	14.774	12.327	11.667	11.236
LSR J1610-0040	14.903	12.911	12.302	12.019